

Measurement of azimuthal anisotropy of light nuclei produced in heavy-ion collisions at RHIC

Rihan Haque
(for the STAR Collaboration)
NISER, Bhubaneswar, India

Outline



- Introduction & motivation
- STAR experiment at RHIC
- Results
- Summary

QCD Workshop
TIFR, India

January 27-30, 2015




Azimuthal anisotropy



Interactions
↓
Pressure (P)

$$y > x \longrightarrow \frac{\partial P}{\partial x} > \frac{\partial P}{\partial y}$$




Azimuthal distribution of produced particles can be described as a Fourier series. The second order coefficient,

$$v_2 = \langle \cos(2(\phi - \psi_R)) \rangle = \left\langle \frac{p_x^2 - p_y^2}{p_x^2 + p_y^2} \right\rangle$$

– Sensitive to early times in the evolution of the system

An estimate of ψ_R , namely Event Plane (ψ_2), is calculated using produced particles in mid-rapidity.



✓ hadron v_2 show constituent quark (NCQ) scaling.

➤ Nuclei are expected to form at a later stage due to their low binding energy

→ Can we expect mass number scaling of nuclei v_2 ?

→ How does nuclei and anti-nuclei v_2 compare?

→ Is there any centrality dependence of nuclei v_2 ?

J. I. Kapusta, Phys. Rev. C 21, 1301 (1980)

R. Scheibl, U. Heinz, Phys. Rev. C 59, 1585 (1999)


D. Molnár, S. A. Voloshin, Phys. Rev. Lett. 91, 92301 (2003)

Figure ref: Phys. Rev. C 75, 054906 (2007), Phys. Rev. Lett. 99, 112301 (2007), Phys. Rev. Lett. 98, 162301 (2007)



The STAR experiment

The Solenoidal Tracker At RHIC (STAR)



Light nuclei identification using TPC

d, dbar, triton: $p_T \sim 1.0$ GeV/c, and ^3He up to 4.5 GeV/c.

Light nuclei identification using ToF

d, dbar, triton: $p_T \sim 4.0$ GeV/c.

1. Time Projection Chamber (TPC)

pseudo-rapidity window: $-1.0 < \eta < 1.0$
full azimuthal coverage.

M Anderson *et al.*, NIM, A 499, 624 (2003)


2. Time of Flight (ToF)

pseudo-rapidity window: $-0.9 < \eta < 0.9$
full azimuthal coverage.

W. J. Llope *et al.*, NIM, B 241, 306 (2005)

Using TPC and ToF π , K , p can be identified up to $p_T \sim 3.0$ GeV/c.


Phys. Rev. C 88, 014902 (2013)



Theoretical dE/dx values: H. Bichsel, NIM, A 562, 154 (2006)




Measurement of nuclei v_2



- ✓ Elliptic flow of d , $\bar{d}$, t , ${}^3\text{He}$, $\bar{{}^3\text{He}}$ measured at mid-rapidity.
- ✓ η sub-event plane method was used with $\eta\text{-gap} = 0.1$
- ✓ some particles not plotted at lower beam energies due to statistical limitations

proton v_2 from Phys. Rev. C 88, 014902 (2013)


Mass ordering of v_2



→ Nuclei v_2 shows mass ordering at low p_T similar to hadrons


hadron v_2 from Phys. Rev. C 88, 014902 (2013)

v_2 of triton (t) and ^3He




$\rightarrow v_2$ of t and ^3He are of similar magnitude (within uncertainty)

v_2 of particles and anti-particles




- Nuclei and anti-nuclei shows similar magnitude of difference in v_2
- The v_2 difference between particle and anti-particle become larger as collision energy decreases, though errors are large at low energies.
- Statistics for anti-nuclei at beam energies below 19.6 GeV is too low to check this consistency.

Centrality dependence of nuclei v_2



- Nuclei v_2 shows centrality dependence for all energies
- antiparticles not shown where statistically limited


Mass number scaling of v_2



Nuclei v_2 show mass number scaling for $p_T / A \sim 1.5 \text{ GeV/c}$ for all beam energies
 $\rightarrow$ Support the general idea that nuclei are formed by coalescence of nucleons

(anti-) proton v_2 from Phys. Rev. C 88, 014902 (2013)

Coalescence model results



Coalescence model agrees with data

→ Another indication of coalescence of nucleons to form nuclei

- ✓ Probability for producing a nucleus is given by the overlap of nucleon phase-space distribution with the Wigner phase-space function of nucleons inside the nuclei.
- ✓ Nucleon phase space information used from a transport (AMPT) model.

R. Mattiello et al. Phys. Rev. Lett. 74, 2180 (1995), L. W. Chen et al. Phys. Rev. C 68, 017601 (2003)
AMPT model: Zi-Wei Lin et al. Phys. Rev. C 72, 064901 (2005)

(A) New Measurement presented:

- ✓ Energy ($\sqrt{s_{NN}} = 7.7, 11.5, 19.6, 27, 39, 62.4$ and 200 GeV) and centrality dependence of nuclei v_2 presented.

(B) Observation and Physics conclusion:

1. Nuclei v_2 versus p_T shows a clear centrality dependence and mass ordering when compared with identified hadrons at all beam energies studied
 - *Mass ordering of v_2 occurs naturally according to hydrodynamic theory of heavy-ion collisions.*
2. Nuclei v_2 versus p_T shows mass number scaling up to $p_T/A = 1.5$ GeV/c
 - *Supports the physics picture of coalescence of nucleons as the dominant mechanism for light nuclei production.*
 - *Results from AMPT+Coalescence calculation are consistent with data.*



Thanks..

Acknowledgements:

STAR Collaboration, NERSC Grid (LBNL), RCAS Grid (BNL),
VECC TIER2 Grid (VECC), KONARK Grid (NISER).

Rihan Haque is supported by DAE-BRNS project grant No. 2010/21/15-BRNS/2026.