


Dark Matter searches at the LHC

Henning Flächer
H.H. Wills Physics Laboratory
University of Bristol

on behalf of the LHC collaborations


SUSY17 –TIFR – Mumbai – 12 Dec 2017

- Overwhelming evidence for Dark Matter but corresponding particle missing from SM
- When searching for Dark Matter, the big question is:
 - How does it interact?
- All evidence for Dark Matter so far is gravitational
 - Why look for it at colliders?



Introduction/Motivation

- Why look for DM at colliders?
- Assumption that in early universe DM was in thermal equilibrium with SM matter
 - some interaction with SM matter
- As universe expands and cools down, DM decouples
- DM abundance determined by annihilation cross section at freeze-out
- A particle with weak scale interactions and mass of $O(100 \text{ GeV})$ gives relic density in agreement with our measurements
 - “WIMP miracle”




$$\Omega_\chi h^2 \simeq 0.1 \times \left(\frac{3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}}{\langle \sigma v \rangle} \right)$$


$$\begin{aligned} \langle \sigma v \rangle &\sim 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1} \\ &\sim \pi \alpha^2 / (100 \text{ GeV})^2 \end{aligned}$$

What interactions to probe?

- Start with minimal assumptions
 - Effective Field Theories



- Simplified models
 - Resolve the interaction





- Described in terms of Lorentz structure, DM mass and cut-off scale
- Need to be careful at LHC @ 13 TeV
- EFT only valid if $Q^2 \ll M$

- Come with different assumptions for interactions/mediators
 - scalar ($\psi\psi$)
 - pseudo scalar ($\psi\gamma^5\psi$)
 - vector $\psi\gamma^\mu\psi$,
 - axial-vector ($\psi\gamma^\mu\gamma^5\psi$)

What interactions to probe?


- Start with minimal assumptions
- Effective Field Theories
- Simplified models
- Resolve the interaction



We also typically need additional radiation (ISR)
as WIMPs are invisible to the detectors

Simplified Dark Matter models

- Typically parameterized by 5 parameters:



- mass of DM particle, m_χ
- mass and width of mediator particle, $m_{\text{med}}, \Gamma_{\text{med}}$
- coupling of mediator to SM sector, g_q
- coupling of mediator to DM sector, g_χ

• A/A-V: $g_q = 0.25, g_\chi = 1$

• S/P-S: $g_q = 1, g_\chi = 1$

- Searches focus on two main signatures:
- DM production via mediator
 - Missing energy searches as a result of escaping DM particles
 - Mono-X signatures
 - $X = \text{jet, photon, } W, Z, H, \dots$
- Resonant production of mediator particle and decay back to SM particles
 - di-jet, di-lepton, di-top, ... resonances

Dark Matter LHC Working Group

- Systematic approach pursued through **Dark Matter LHC Working group**
 - <https://lpcc.web.cern.ch/content/lhc-dm-wg-dark-matter-searches-lhc>
 - with involvement of both experimentalists and theorists
- Recommendations for models and their implementation
- Guidelines on how to compare collider searches with direct detection limits
- Recommendations for comparison of searches for heavy mediators of DM production

Dark Matter Benchmark Models for Early LHC Run-2 Searches:
Report of the ATLAS/CMS Dark Matter Forum

August 8, 2016

Daniel Abercrombie *MIT, USA*

Nural Akchurin *Texas Tech University, USA*

Ece Akilli *Université de C*

Juan Alcaraz Maestre *C*

(CIEMAT), Spain

Brandon Allen *MIT, USA*

Barbara Alvarez Gonzale

Jeremy Andrea *Institut P*

Université de Strasbourg

Alexandre Arbey *Univers*

École Normale Supérieu

Georges Azuelos *Univers*

Patrizia Azzi *INFN Padov*

Mihailo Backović *Centre*

catholique de Louvain, B

Yang Bai *Department of*

Swagato Banerjee *Unive*

James Beacham *Ohio St*

Alexander Belyaev *Ruthe*

dom

Antonio Boveia *(editor) C*

Amelia Jean Brennan *Th*

Oliver Buchmueller *Impe*

Matthew R. Buckley *Dep*

Giorgio Busoni *SISSA ar*

Michael Buttiglione *Institut*

Université de Strasbourg

Giacomo Cacciapaglia *U*

France

Regina Caputo *Santa Cr*

of Astronomy and Astrop

Linda Carpenter *Ohio St*

Nuno Filipe Castro *LIP-N*

Ciências da Universidade

Guillermo Gomez Cebal

Yangyang Cheng *Univers*

John Paul Chou *Rutgers*

Arely Cortes Gonzalez *If*

Editor

¹CERN, EP Department, CH-1211

²High Energy Physics Group, Black

London, SW7 2AZ, United Kingd

³ARC Centre of Excellence for Pa

University of Melbourne, 3010, Aust

⁴UC, Santa Cruz and UC, Santa C

⁵Antwerp University, B2610 Wilrij

⁶SISSA and INFN Sezione di Tri

⁷Fysiska institutionen, Lunds univ

⁸LPSC, Université Grenoble-Alpes

⁹Department of Physics and Astron

USA

¹⁰Rudolf Peierls Centre for Theoret

United Kingdom

¹¹Department of Physics and Astron

USA

¹²Department of Physics and Astron

USA

¹³Department of Physics and Astron

USA

¹⁴Department of Physics and Astron

USA

¹⁵Department of Physics and Astron

USA

¹⁶Department of Physics and Astron

USA

¹⁷Department of Physics and Astron

USA

¹⁸Department of Physics and Astron

USA

¹⁹Department of Physics and Astron

USA

²⁰Department of Physics and Astron

USA

²¹Department of Physics and Astron

USA

²²Department of Physics and Astron

USA

²³Department of Physics and Astron

USA

²⁴Department of Physics and Astron

USA

²⁵Department of Physics and Astron

USA

²⁶Department of Physics and Astron

USA

²⁷Department of Physics and Astron

USA

²⁸Department of Physics and Astron

USA

²⁹Department of Physics and Astron

USA

³⁰Department of Physics and Astron

USA

³¹Department of Physics and Astron

USA

³²Department of Physics and Astron

USA

³³Department of Physics and Astron

USA

³⁴Department of Physics and Astron

USA

³⁵Department of Physics and Astron

USA

³⁶Department of Physics and Astron

USA

³⁷Department of Physics and Astron

USA

³⁸Department of Physics and Astron

USA

³⁹Department of Physics and Astron

USA

⁴⁰Department of Physics and Astron

USA

⁴¹Department of Physics and Astron

USA

⁴²Department of Physics and Astron

USA

⁴³Department of Physics and Astron

USA

⁴⁴Department of Physics and Astron

USA

⁴⁵Department of Physics and Astron

USA

⁴⁶Department of Physics and Astron

USA

⁴⁷Department of Physics and Astron

USA

⁴⁸Department of Physics and Astron

USA

⁴⁹Department of Physics and Astron

USA

⁵⁰Department of Physics and Astron

USA

⁵¹Department of Physics and Astron

USA

⁵²Department of Physics and Astron

USA

⁵³Department of Physics and Astron

USA

⁵⁴Department of Physics and Astron

USA

⁵⁵Department of Physics and Astron

USA

⁵⁶Department of Physics and Astron

USA

⁵⁷Department of Physics and Astron

USA

⁵⁸Department of Physics and Astron

USA

⁵⁹Department of Physics and Astron

USA

⁶⁰Department of Physics and Astron

USA

⁶¹Department of Physics and Astron

USA

⁶²Department of Physics and Astron

USA

⁶³Department of Physics and Astron

USA

⁶⁴Department of Physics and Astron

USA

⁶⁵Department of Physics and Astron

USA

⁶⁶Department of Physics and Astron

USA

⁶⁷Department of Physics and Astron

USA

⁶⁸Department of Physics and Astron

USA

⁶⁹Department of Physics and Astron

USA

⁷⁰Department of Physics and Astron

USA

⁷¹Department of Physics and Astron

USA

⁷²Department of Physics and Astron

USA

⁷³Department of Physics and Astron

USA

⁷⁴Department of Physics and Astron

USA

⁷⁵Department of Physics and Astron

USA

⁷⁶Department of Physics and Astron

USA

⁷⁷Department of Physics and Astron

USA

⁷⁸Department of Physics and Astron

USA

⁷⁹Department of Physics and Astron

USA

⁸⁰Department of Physics and Astron

USA

⁸¹Department of Physics and Astron

USA

⁸²Department of Physics and Astron

USA

⁸³Department of Physics and Astron

USA

⁸⁴Department of Physics and Astron

USA

⁸⁵Department of Physics and Astron

USA

⁸⁶Department of Physics and Astron

USA

⁸⁷Department of Physics and Astron

USA

⁸⁸Department of Physics and Astron

USA

⁸⁹Department of Physics and Astron

USA

⁹⁰Department of Physics and Astron

USA

⁹¹Department of Physics and Astron

USA

⁹²Department of Physics and Astron

USA

⁹³Department of Physics and Astron

USA

⁹⁴Department of Physics and Astron

USA

⁹⁵Department of Physics and Astron

USA

⁹⁶Department of Physics and Astron

USA

⁹⁷Department of Physics and Astron

USA

⁹⁸Department of Physics and Astron

USA

⁹⁹Department of Physics and Astron

USA

¹⁰⁰Department of Physics and Astron

USA

¹⁰¹Department of Physics and Astron

USA

¹⁰²Department of Physics and Astron

USA

¹⁰³Department of Physics and Astron

USA

¹⁰⁴Department of Physics and Astron

USA

¹⁰⁵Department of Physics and Astron

USA

¹⁰⁶Department of Physics and Astron

USA

¹⁰⁷Department of Physics and Astron

USA

¹⁰⁸Department of Physics and Astron

USA

¹⁰⁹Department of Physics and Astron



USA

¹¹⁰Department of


Simplified Dark Matter models

Mono-mania @ the LHC...


Mono-Z




Mono-Higgs



Mono-W



Mono-top





... complemented by searches for “full”, UV-complete models, such as

- Supersymmetry
- Kaluza-Klein extra dimensions
- Little Higgs models
- etc.


- ...and of course there are also other DM candidates but this talk will mainly focus on WIMPs

Mono-mania





Mono-jet & Mono-V Searches

- Search for jets with missing energy
 - Jets or boosted vector bosons (W, Z)
- Comparison of data with background prediction
- Main background from $Z \rightarrow vv + \text{jets}$ and $W+\text{jets}$ production




CMS EXO-16-048







ATLAS EXOT-2016-27

Mono-jet & Mono-V Searches

- Interpretation in simplified DM models
- s-channel with different mediators





- Set limits on allowed masses and coupling strength (next slide)
- Further interpretations in
 - fermion portal models
 - coloured scalar mediator models
 - nonthermal dark matter model



Mono-jet & Mono-V Searches


- Interpretation in simplified DM models
- s-channel with different mediators




CMS EXO-16-048

Mono-photon

- Photon $p_T > 150$ GeV
- Veto events with more than 1 jet
- Require angular separation between photon and missing energy





ATLAS:
[Eur. Phys. J. C 77 \(2017\) 393](https://doi.org/10.1140/epjc/s10050-017-4880-0)



Weaker sensitivity
 compared to
 mono-jet ISR
 searches




But also EFTs with
 direct couplings to
 photons

Comparable CMS analysis:
[JHEP 10 \(2017\) 073](https://doi.org/10.1007/JHEP10(2017)073)




Mono-Z

- Search for dark matter candidates produced in association with a Z boson
- [ATLAS: PLB 776 \(2017\) 318, arXiv:1708.09624](#)





Vector



Axial-Vector

CMS-EXO-16-052




Scalar almost identical

Pseudo-
scalar


Higgs as a portal to Dark Matter

- Study of Higgs \rightarrow invisible branching fraction in **VBF**, association with V and gluon fusion production modes.
- As Higgs couples proportionally to mass, making it a viable portal to DM
- Only sensitive to DM masses $< 0.5 m_H$




- Additional searches for Higgs in conjunction with DM in bb and $\gamma\gamma$ final states (\rightarrow backup)

Higgs production in association with $Z \rightarrow ll$
CMS-EXO-16-052





- Combination of $H \rightarrow$ invisible production modes
- $B(H \rightarrow \text{inv}) < 0.24$ @95%CL

CMS-HIG-16-016




DM in association with heavy flavours

- DM in association with bottom or top quarks
- VBF like signature, b-jet tagging



- Interpretation in colour-neutral $t\bar{t}/bb + \phi$ scalar and $t\bar{t}/bb + a$ pseudoscalar models
- Not yet sensitive to $bb + \phi/a$ production
- Also searches with single top
- CMS EXO-16-051




arXiv:1710.11412


- Fixed DM mass
- Limit as fcn of mediator mass

Scalar



Similar sensitivity to pseudoscalar



- Fixed mediator mass
- Limit as fcn of DM mass





Mediator Searches



High-mass dijet resonances


- Search for resonance in dijet invariant mass spectrum




Low-mass dijet resonances

- Low mass mediators are difficult to constrain because of huge QCD dijet background
 - difficult/impossible to cope with total event rate
- Possible ways out:
 - “**Data scouting**”: perform analysis on dataset that contains reduced event information (trigger level), allowing to store data at very high rate
 - Trigger on **high-pT ISR jet or photon** and search for low mass resonance in recoil system
 - Trigger on high-pT ISR jet and search for merged, boosted resonance in recoil, using **jet substructure**

ATLAS Trigger Level Analysis ATLAS-CONF-2016-030




ATLAS ISR search (photon) ATLAS-CONF-2016-070




Low-mass dijets— boosted topology

- Resonance is produced with sufficiently high transverse momentum that its decay products are merged into a single jet with two-prong substructure.






CMS-EXO-17-001

- Soft drop mass for jets in different p_T ranges
- Peak from Ws and Zs clearly visible
- Sensitive to dijet resonances with masses as low as 50 GeV



Dilepton resonances (Z')


- $Z' \rightarrow e^+e^-$ and $Z' \rightarrow \mu^+\mu^-$ searches
 - Require high- p_T same flavour, opposite charge dilepton pair
 - Limits on Z' mass of around 4 - 4.5 TeV



Model	Width [%]	θ_{E_6} [rad]	Lower limits on $M_{Z'}$ [TeV]					
			ee		$\mu\mu$		$\ell\ell$	
			Obs	Exp	Obs	Exp	Obs	Exp
Z'_{SSM}	3.0	-	4.3	4.3	4.0	3.9	4.5	4.5
Z'_{χ}	1.2	0.50π	3.9	3.9	3.6	3.6	4.1	4.0
Z'_{S}	1.2	0.63π	3.9	3.8	3.6	3.5	4.0	4.0
Z'_{I}	1.1	0.71π	3.8	3.8	3.5	3.4	4.0	3.9
Z'_{η}	0.6	0.21π	3.7	3.7	3.4	3.3	3.9	3.8
Z'_{N}	0.6	-0.08π	3.6	3.6	3.4	3.3	3.8	3.8
Z'_{ψ}	0.5	0π	3.6	3.6	3.3	3.2	3.8	3.7


The grand picture

- Can combine searches for DM signal (missing energy) and those for mediators to constrain allowed model parameter space




The grand picture

- Can combine searches for DM signal (missing energy) and those for mediators to constrain allowed model parameter space
- Exclusion strongly depends on choices for additional parameters, e.g., coupling to quarks but also leptons!




The grand picture

- Can combine searches for DM signal (missing energy) and those for mediators to constrain allowed model parameter space
- Exclusion strongly depends on choices for additional parameters, e.g., coupling to quarks but also leptons!





The grand picture

- Can combine searches for DM signal (missing energy) and those for mediators to constrain allowed model parameter space
- Exclusion strongly depends on choices for additional parameters, e.g., coupling to quarks but also leptons!



The grand picture


- Much weaker constraints on Scalar and Pseudo-Scalar mediators
- Exclusion only for very light DM and couplings $g_q = g_{DM} = 1$.




Comparison with Direct Detection

- Complementarity with Direct Detection clearly visible
- Sensitivity strongly depends on type of DM interaction
 - Can be used to learn about underlying physics in case one or the other sees a signal
- Collider searches more powerful for small DM masses

Spin-independent




Spin-dependent



Comparison with Direct Detection

- Constraints on $B(H \rightarrow \text{inv})$ can be translated to limits in scattering cross section vs DM mass plane to compare with direct detection experiments.
- Sensitive to masses $< 0.5 m_H$
- very sensitive at low DM masses




Wimp – Nucleon Interaction	
<u>Spin-Independent (SI)</u>	<u>Spin-Dependent (SD)</u>
Basic Mediators	
Vector <i>Besides low DM masses DD provides best sensitivity. Complementarity at low DM masses (<5 GeV)!</i>	Axial-vector <i>DD and collider are equal in overall sensitivity but probe different regions of parameter space! Complementarity in full parameter space!</i>
Scalar <i>Besides low DM masses DD provides best sensitivity. Complementarity at low DM masses (<5 GeV)!</i>	Pseudoscalar <i>Effectively no limits from DD above a few GeV in M_{med}. Collider and ID probe region at larger M_{med}. Complementarity in M_{med}!</i>

from
O. Buchmueller

Outlook – Where next?

Axial-Vector mediator


$$g_q = g_\gamma = 1$$



from arXiv:1409.4075, Phys.Dark Univ. 9-10 (2015) 51-58




Vector mediator

$$g_q = g_\gamma = 1$$



Dark Photon search @ LHCb

- Assume a dark sector, a collection of particles that are not charged directly under the SM strong, weak, or electromagnetic forces.
- Dark photon, A' , whose coupling to the electromagnetic current is suppressed relative to that of the ordinary photon, γ , by a factor of ε
- Dark Photon would also couple to DM
- Search for Dark Photon decay to di-muon pair in meson decays





Summary

- Comprehensive search programme for dark matter is underway at LHC
- So far focused on WIMP signatures
 - Guidance from simplified Dark Matter models
 - Search strategy includes searching for DM production but also for mediators that couple to both SM and DM
- **Long-lived signatures** to be added to search portfolio
- Extend **searches beyond WIMP** signatures
- DM searches @ LHC a powerful tool
 - In particular sensitivity to small DM masses
 - **Sensitivity strongly depends on assumptions** made for DM interaction, spin structure of mediator and coupling strength
 - In many ways **complementary to direct detection** searches
 - This could be crucial for understanding the underlying physics in case of a signal from either approach



- **Backup**

Mono-jet & Mono-V Searches


- Interpretation in simplified DM models
- Fermion portal models





- Interpretation in simplified DM models
- Coloured scalar mediator models



- Interpretation in colour-neutral $t\bar{t}/bb\bar{b} + \phi$ scalar and $t\bar{t}/bb\bar{b} + a$ pseudoscalar models
 - Fixed DM mass
 - Limit as fcn of mediator mass
 - Fixed mediator mass
 - Limit as fcn of DM mass
- Not yet sensitive to $bb + \phi/a$ production





- colour-charged scalar mediators (b-FDM)





Mono-Higgs

- Higgs production in conjunction with DM
 - Higgs decay to $\gamma\gamma$



CMS-PAS-EXO-16-054


- Higgs production in conjunction with DM
 - Higgs decay to bb



[Phys. Rev. Lett. 119 \(2017\) 181804](https://doi.org/10.1103/PhysRevLett.119.181804)


High-mass dijet resonances

- Search for resonance in dijet invariant mass spectrum



- "bump hunt" not effective for very wide resonances
- Investigate di-jet scattering angle in this case

DM mediator
would show
up here



Comparison with Direct Detection

- Complementarity with Direct Detection clearly visible
- Sensitivity strongly depends on type of DM interaction
 - Can be used to learn about underlying physics in case one or the other sees a signal
- Collider searches more powerful for small DM masses

