


High energy neutrinos


Danny Marfatia


Cosmic neutrinos at IceCube



- 50 events with deposited energy > 60 TeV
- 3 cascade events between 1-2 PeV


Galactic or extragalactic?






- Galactic fraction < 9.5% at 90% CL
- Zero galactic flux allowed at < 1 sigma


Same source for UHECR and cosmic neutrinos?






Consistent with the Waxman-Bahcall flux

Upgoing muon neutrinos




1710.01191



Upgoing muon neutrino spectrum ($E > 120 \text{ TeV}$)
harder than HESE spectrum

Multicomponent flux?




Glashow resonance not observed


$\bar{\nu}_e$ is unique because of resonant scattering at

$$E_\nu = \frac{M_W^2}{2m_e} = 6.3 \text{ PeV}$$

$\bar{\nu}_e e^- \rightarrow W^- \rightarrow \text{anything}$



Connection with gamma rays



- Neutrino spectra softer than shown are inconsistent with Fermi data
- Connection for $p\gamma$ sources weaker because target photons prevent gamma rays from leaving source

Lorentz and CPT violation?

with Jiajun Liao (1711.09266)

- Suppose LIV and CPTV only occur in neutrino sector
- Only consider effects that change the kinematics of particle interactions
- Postulate that CPTV arises from Planck-suppressed terms in the Lagrangian


- Modified dispersion relation

$$E^2 - p^2 = m^2 + 2\delta E^2$$

$$\delta = \kappa \frac{E}{M_{Pl}}$$

- Assume all neutrino flavors have same LIV parameter to be consistent with neutrino oscillation data
- Dispersion relation for antineutrinos: $\delta \rightarrow -\delta$
- Our choice $\delta > 0 \Rightarrow$ neutrinos are superluminal and antineutrinos are subluminal

- Dominant energy loss processes for superluminal neutrinos are vacuum pair emission (VPE) and neutrino splitting



- **Event pile-up** caused by neutrino splitting is larger than for VPE because splitting produces 2 additional lower energy neutrinos

$$\Gamma \propto \kappa^3 \frac{G_F^2 E^8}{M_{Pl}^3}$$

Effect on neutrino sources?

$\pi^+ \rightarrow \mu^+ \nu_\mu$ requires the energy of superluminal neutrinos to satisfy

$$E^3 \leq \frac{(m_\pi - m_\mu)^2 M_{Pl}}{2\kappa}$$

VPE occurs above an energy threshold given by

$$E_{th}^3 = \frac{2m_e^2 M_{Pl}}{\kappa}$$

Threshold energy for neutrino splitting is tiny compared to that for VPE

For a given VPE threshold energy, the upper bound on the superluminal neutrino energy is

$$E < 10.3E_{th}$$

If the highest energy track event observed by IceCube (with median estimated energy of 8.7 PeV) was initiated by a superluminal neutrino, then

$$E_{th}^{min} = 0.85 \text{ PeV}$$

Need extragalactic sources of superluminal neutrinos

CR Reservoirs:

Starburst galaxies and Galaxy clusters

$$pp \rightarrow \pi^\pm \text{ pairs} \rightarrow \nu_e + \bar{\nu}_e + 2\nu_\mu + 2\bar{\nu}_\mu$$

- naturally produces spectral break
- same number of neutrinos and antineutrinos

CR Accelerators:

Gamma-ray bursts and Active Galactic Nuclei

$$p\gamma \rightarrow \pi^+ \rightarrow \nu_e + \nu_\mu + \bar{\nu}_\mu$$

- naturally produces hard neutrino spectrum
- twice as many neutrinos and antineutrinos

If muons damped

$$p\gamma \rightarrow \pi^+ \rightarrow \nu_\mu \text{ only}$$

Only superluminal neutrinos at source!

BUT, intrinsic contamination from π^- is expected to reduce the superluminal fraction by 20%-33%


e.g.

$$p\gamma \rightarrow n\pi^+$$


$$n\gamma \rightarrow p\pi^-$$

Flavor ratio at Earth


	Source flavor ratio	Earthly flavor ratio	$\bar{\nu}_e$ fraction in flux (\mathcal{R})		
$pp \rightarrow \pi^\pm$ pairs	(1:2:0)	(1:1:1)	18/108 = 0.17		
w/ damped μ^\pm	(0:1:0)	(4:7:7)	12/108 = 0.11		
$p\gamma \rightarrow \pi^+$ only	(1:1:0)	(0:1:0)	(14:11:11)	(4:7:7)	8/108 = 0.074
w/ damped μ^+	(0:1:0)	(0:0:0)	(4:7:7)	(0:0:0)	0
charm decay	(1:1:0)	(14:11:11)	21/108 = 0.19		
neutron decay	(0:0:0)	(1:0:0)	(0:0:0)	(5:2:2)	60/108 = 0.56




No asymmetry in neutrino-antineutrino composition



	$E^{-2.92}(1:1:1)$	$E^{-2}(1:1:1)$	E^{-2} with CPTV		
Case	(a)	(b)	(a)	(b)	(a)
χ^2	9.6	10.3	24.0	34.0	7.7
GR events	0.16	0.15	3.1	2.7	0.98



Superluminal fraction compatible with π^- contamination



1711.09266

Summary

- Multicomponent neutrino flux not required if neutrino interactions violate CPT
- Single E^{-2} spectrum from muon-damped $p\gamma$ source works
- Superluminal fraction compatible with π^- contamination
- Excess below 200 TeV explained by event pile-up from superluminal neutrino decay
- Subluminal antineutrinos contribute at high energies so no cutoff in spectrum
- Expect Glashow resonance events soon